



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

## NOTAE MYCOLOGICAE, SER. XXIX

### Micromycetes Dakotenses et Utahenses a Doct. J. F. Brenckle Lecti et Communicati

P. A. SACCARDO

#### A. TELEOMYCETAE

1. *ROSELLINIA AMPHISPHAERIOIDES* Sacc. et Speg., Syll. I: 262, F. Ital. fig. 588

Hab. in ramis putrescentibus *Ribis floridani*, Kulm, N. D., Oct., 1916 (1205).

Asci  $100 \times 7-8$ , stip.  $20-26 \mu$  longis; sporidia 1-sticha  $17-21 \times 7-8$ , fuliginea, senio longitrosum rimosa ut in *Anthostomate taeniosporo*. Esse potest forma, a loco sub cortice relaxato, deminuta et deformata, *R. mastoideae* Sacc.

2. *Rosellinia subsimilis* Sacc., sp. nov.

Péritheciis gregariis, subglobosis, minutis, nigris, glabris,  $100-180 \mu$  diam. brevissime papillatis; ascis filiformi-paraphysatis, cylindricis, subsessilibus,  $89-90 \times 5.5-6$ , 8-sporis; sporidiis oblique monostichis e cylindraceo ellipsoideo-oblongis, saepius leviter inaequilateris, utrinque rotundatis,  $12-14 \times 5-6$ , fuligineis.

Hab. in ramis decorticates emortuis *Crataegi* sp. Whitestone Gully,<sup>1</sup> N. D., Nov., 1916 (1188). Affinis *R. rimincolae* differt sporidiis angustioribus subcylindraceis, etc.

3. *MELANOPSAMMA POMIFORMIS* (Pers.) Sacc., Syll. I: 575

Hab. in ramis putrescentibus *Corni stoloniferae*. Whitestone Gully,<sup>1</sup> N. D., Nov., 1916 (1185). Specimina immatura et non omnino certa.

4. *DIDYMELLA EUPYRENE* Sacc., Syll. I: 554

Hab. in caulibus putrescentibus *Urticae gracilis*, Ft. Douglas, Utah, Majo, 1918 (53).

<sup>1</sup> Whitestone Gully extends from and lies east of the U. S. Monument of "Whitestone Battlefield" in Dickey County.

5. *DIDYMELLA EFFUSA* (Niessl) Sacc., Syll. **I**: 552

Hab. in caule emortuo *Artemisiae biennis*, Kulm, N. D., Sept., 1914 (632). Forma ascis 60–70  $\times$  7, sporidiis breviter fusoidis, 12–15  $\times$  4, plasmate bipartito 1-septatis, vix constrictis.

6. *MELANOMMA MEDIUM* Sacc. et Speg., Syll. **II**: 104

Hab. in ramis decorticatis putrescentibus *Sambuci glaucae*, Ft. Douglas, Utah, Apr., 1918 (70).

Asci 90–100  $\times$  14; sporidia disticha v. oblique 1-sticha, teretio-oblunga, utrinque rotundata, 22–25  $\times$  7–8, 3-septata, leviter constricta, dilute brunnea.

7. *TEICHOSPORA SOLITARIA* Ellis, Pyr. N. Amer., p. 214, *Cucurbitaria* Ell. Syll. **II**: 321.

Hab. in trunco decorticato et intemperie indurato *Atriplicis confertifoliae*, Grantville, Utah, Apr., 1918 (32).

Valde affinis est *T. oxythele* Sacc. et Briand. Syll. **IX**: 903. Berl. *Jc. II*, t. 74.

8. *PLEOSPORA SHEPHERDIAE* Peck., Syll. **IX**: 876

Hab. in ramis corticatis emortuis *Shepherdiae argenteae*, Whitestone Gully, N. D., Nov., 1918 (1192).

9. *HYPOXYLON EFFUSUM* Nits., Syll. **I**: 379

Hab. ad lignis putrescentia, Fort Riley, Kans., Oct., 1917 (101).

**Phaeotrype** Sacc., gen. nov. (Etym. quasi *Diatrype phaeospora*)

Stromata pulvinata, erumpenti superficialia, nigra, ostiolis vix exantibus punctulata; perithecia pauca monosticha, substantia discolori excepta. Asci breviter stipitati, octospori, aparaphysati (?). Sporidia allantoidea, majuscula, atro-olivacea. Est omnino *Diatrype* sed vere phaeospora.

10. **Phaeotrype Brencklei** Sacc., sp. nov.

Stromatibus gregariis, subsuperficialibus, peridermii laciniis basi cinctis, depresso pulvinatis, atro-nitidulis, 0.5  $\times$  0.7 mm.

diam., superficie levissime colliculosa et punctata; intus sordide albido, carnosulo; peritheciis 8-10 monostichis, globulosis, 0.3 mm. diam., ostiolis punctiformis, obtusis, integris; ascis fusoides breviter stipitatis,  $80-85 \times 8-9$ , octosporis; sporidiis distichis allantoideis curvis, utrinque rotundatis;  $16-17 \times 3.5-4$ , atro-olivaceis, subnigricantibus.

Hab. in ramis corticatis emortuis *Rosae* sp., Whitestone Gully, N. Dak., Nov., 1916 (1198).

11. *Diatrype paurospora* Sacc., sp. nov.

Stromatibus gregariis, erumpenti-superficialibus, peridermo cinctis, pulvinatis, nigris, duris, fragilibus, superficie colliculosis v. rimosis, 1-2 mm. diam.; peritheciis paucis, globosis, 0.3 mm diam., substantia sorde grisea exceptis; ostiolis vix excedentibus obtusis; ascis fusoides, apice rotundatis,  $35-40 \times 7$ , subsessilibus, aparaphysatis (?), 2-4-sporis; sporidiis allantoideis, curvulis, utrinque rotundatis,  $16 \times 4-5$ , hyalinis, vix chlorinis, tunica crassiuscula.

Hab. in ramis corticatis emortuis *Quercus utahensis* pr. Ft. Douglas, Utah, Jan., 1918 (12). Imprimis ascis tantum 2-4-sporis distinguenda species.

12. *EUTYPA SCABROSA* (Bull.) Fuck., Syll. I: 171

Hab. in ramis emortuis corticatis et decorticatis *Amelanchieris alnifoliae*, Whitestone Gully, Nov., 1916 (1199).

13. *EUTYPA LEIOPLACA* (Fr.) Cooke, Syll. I: 170

Hab. in ramis decorticatis emortuis *Amelanchieris alnifoliae*, Whitestone Gully, Nov., 1916 (1182). Socia adest forma diminuta *Calosphaeriae Principis*. Specimina *Eutypae* juvenilia et tunc dubium an potius *Eutypam latam* spectent.

14. *EUTYPELLA SARCOBATI* Ellis et Ev., Syll. XIV: 484

Hab. in ramis corticatis *Sarcobati vermiculati*, Salt Lake pr. Grantville, Utah, Apr., 1918 (29). Superficiis ligni late mycelio atrata, hinc speciis ad *Eutypam* nutat.

15. *VALSA CORNICOLA* Cooke, Syll. I: 122

Hab. in ramis corticatis emortuis *Corni stoloniferae*, Whitestone Gully, N. D., Nov., 1916 (1184). Est forma minor.

16. *VALSA CINCTA* Fr., Syll. I: 142

Hab. in ramis corticatis *Amelanchieris*?, Whitestone Gully, Nov., 1916 (1183a).

17. *VALSA LEUCOSTOMA* (Pers.) Fr., Syll. I: 139

Hab. in ramis corticatis *Pruni melanocarpi*, Ft. Douglas, Utah, Jul., 1918 (85). Specimina non bene evoluta.

18. *Chorostate utahensis* Sacc., sp. nov.

Stromatibus erumpentibus, prominulis pulvinatis, 1.5 mm. diam., nigricantibus; peritheciis numerosis, monostichis, 0.3 mm. diam., globosis, ostioli punctiformibus, vix extantibus; ascis cylindraceis apice rotundatis, ibique non v. vix lumine bifoveolatis, breviter stipitatis,  $55-60 \times 7.3-8$ , apapophysatis, octosporis; sporidiis oblique monostichis, oblongo-clavatis, curvatis,  $12-14 \times 3-6$ , eguttulatis, hyalinis, constricto-1-septatis.

Hab. in ramis corticatis emortuis *Quercus utahensis*, Ft. Douglas, Utah, Mayo, 1918 (94). Affinis *C. teiphaemae*, differt sporidiis curvis, loculo super crassiore rotundato, ascis angustioribus, ostioli abbreviatis.

19. *Diaportha (Euportha) Brenckleana* Sacc., sp. nov.

Peritheciis laxe gregariis, ligno immutato sed linea stromatica percurso immersis, globosis,  $350-500 \mu$  diam., contextu minute celluloso atro-olivaceo; ostioli erumpentibus tereti-conicis parum extantibus; ascis fusoides, apice rotundatis,  $45-55 \times 8-9$ , apapophysatis, octosporis; sporidiis oblique 1-stichis, cylindraceo-fusoides, utrinque rotundatis,  $14 \times 3.6$ , inaequaliter 4-guttulatis, hyalinis, leviter constrictis.

Hab. in ramis corticatis, interdum superficie atratis, *Corni stoloniferae*, Whitestone Gully, N. D., Nov., 1916 (1186). A *D. crassicoli* ostiolo omino diverso, etc. secedit.

## 20. FENESTELLA MINOR Tul., Syll. II: 326

Hab. in ramis corticatis emortuis *Corni stoloniferae*, White-stone Gully, N. D., Nov., 1916 (1194). Asci 100-120  $\times$  10-12, p. spor. 80-85  $\mu$  longa, plerumque 4-spori; sporidia 6-10-septato-muralia, 25-30  $\times$  9-11, fuliginea; stromata valsea.

21. *Lachnum crystalligerum* Sacc., sp. nov. (mox endenda)

Hab. in ramis corticatis emortuis *Rubi parviflori*, Ft. Douglas, Utah, Jul., 1918 (81). Specimina non omnino matura.

## 22. PYRENOPEZIZA RUBI (Fr.) Rehm., Syll. VIII: 361

Hab. in ramis emortuis *Rubi strigosi*, Kulm, N. D., Sept., 1916 (1190).

## 23. ORBILIA VINOSA (A. et S.) P. Karst., Syll. VIII: 622

Hab. in lignis putridis pr. Anselm, N. D., Aug., 1916 (1072).

## 24. CENANGIUM FURFURACEUM (Roth.) De Not., Syll. VIII: 565

Hab. in ramis corticatis emortuis *Ceanothi velutini*, Ft. Douglas, Utah, Apr., 1918 (39). Probabiliter haec species, sed specimina imperfecta et tunc paullulum dubia.

25. *Patinella Brenckleana* Sacc., sp. nov.

Ascomatibus gregariis, subsuperficialibus, scutellatis, applanatis, 0.7-1 mm. diam., nigris, glabris, margine rectiusculo, acuto, disco plano intus flavido, epithecio vero grumoso, atro-fulvo; ascis tereti-clavatis, apice rotundatis, deorsum sensim tenuato-stipitatis, 70  $\times$  11-12, octosporis; sporidis distichis, fusoides, saepe leniter curvis, utrinque obtusatis, dilutissime flavidis, farctis; paraphysibus filiformibus, hyalinis.

Hab. in cortice *Amelanchieris alnifoliae*, Whitestone Gully, N. D., Nov., 1916 (1196). Affinis *P. inquinanti* a qua differt ascomate 1 mm. lato, disco plano, sporidiis paullo minoribus, paraphysibus aequalibus.

## B. DEUTEROMYCETAE

26. *SPHAERONAEMA POLYMORPHUM* Auersw., Syll. **III**: 185

Hab. in ramis corticatis *Pruni melanocarphae*, Ft. Douglas, Utah, Nov., 1917 (4). Dicitur status pycnid. *Cenangellae vernicosae* (Fuck.) Sacc.

27. *DIPLODIA ABROTANI* Fuck., Syll. Syll. **III**: 369

Hab. in caulibus emortuis *Artemisiae abrotani*, cultae, Kulm, N. D., Jan., 1917 (1207). Sporulae ovato-ellipsoideae,  $20-24 \times 11$ , diu continuae (ut *Sphaeropsis*) denique obsolete 1-septatae, sporophora interdum in pseudoparaphyses  $20-30 \times 1.5-2$  hyalinas abeunt.

28. *Septoria Lunelliana* Sacc., sp. nov.

Pycnidiis gregariis globoso-depressis, epidermide initio velatis,  $180 \mu$  diam. subastomis, dein late dehiscentibus et collabescentibus, extus nigricantibus; contextu fulvo-brunneo, celluloso; sporulis filiformibus, rectis v. curvulis, utrinque acutulis,  $65-75 \times 2.3-2.5$ , hyalinis, obsolete-3-septatis, guttulatisque.

Hab. in foliis languidis expallentibus *Caricis atrostachyae* pr. Leeds, N. D., Jul., 1915 (J. Lunell, n. 1206). A *Septoriis* cariciolis praecipue pycnidiis contextu fulvescentibus, collabescentibus, sporulis aequalibus, obsolete 3-septatis dignoscitur.

29. *Melanconium botryosum* Sacc., sp. nov.

Acervulis erumpentibus, gregariis, nigris, peridermio laciniato cinctis, ob conidia expulsa et compacta superficie colliculoso-botryosis (sphaeriae-formibus); conidiis copiosissimis, globosis. levibus,  $7-8 \mu$  diam., ochraceo-fuligineis; conidiophoris papilliformibus, exiguis ex hypostromate celluloso, ochraeo-fusco orientibus.

Hab. in ramis corticatis emortuis *Pruni melanocarphae*, Ft. Douglas, Utah, Nov., 1917 (3). Typus singularis et vix *Melanconio* proximus, probibiler gen. nov.

30. *Steganosporium utahense* Sacc., sp. nov.

Acervulis gregariis, subcutaneo-erumpentibus, sed non praeminentibus, 1 mm. diam., primo peridermio bullato tectis, dein ab

ejus laciniis pallidis cinctis, nigricantibus, subceraceis; conidiis polymorphis e globoso-angulosis, v. triangulis, v. gibbis,  $24-31\ \mu$  diam., ex cellulis 3-10, botryoideo-conjunctis formatis, rufo-fulgineis; conidiophoris paliformibus saepe 1-septatis, subhyalinis,  $28 \times 8-9$ , ex hypostromate crasso distincte celluloso rufo-fusco oriundis.

Hab. in ramulis corticatis emortuis *Chrysosporium nauseosi*, Ft. Douglas, Utah, Apr., 1918 (27). Affine *S. heterospermo* Vesterg sed acervulis initio epidermide albida pustulata tectis, dein cinctis, conidiis non foedantibus, conidiophoris distinctis paliformibus, etc. dignoscitur.

31. DENDRYPHIUM NODULOSUM Sacc., Syll. IV: 490.

Hab. in caulibus emortuis putridis *Urticae gracilis*, Nyland's Grove, Lamoure County, N. D. (1181). Probabiliter *Helminthosporium interseminatum* B. et Br. vix differt.

32. EPOCHNIUM ISTHMOPHORUM Sacc., sp. nov. (mox endenda)

Hab. in caulibus emortuis *Chrysosporium nauseosi*, Ft. Douglas, Utah, Apr., 1918 (34). Genus *Epochnium* non tantum ad *Diplococcium* nutat sed et forte propius ad *Bisporam*. Conidia isthmo saepius connexa, usque ad  $20-24 \times 11-12$ .

33. CYLINDROCOLLA URTICAE (Pers.) Bon., Syll. IV: 674.

Hab. in caulibus *Urticae* emortuis, Ft. Douglas, Utah, Majo, 1918 (54).

VIA LUCA BELLUDI, 15,  
PADOVA, ITALY.